

**Appendix D.** Codes and descriptions of 89 SWReGAP land cover types that occur in New Mexico (NatureServe 2004).

Code	Description
S002	<i>Rocky Mountain Alpine Bedrock and Scree</i> - This ecological system is restricted to the highest elevations of the Rocky Mountains, from Alberta and British Columbia south into New Mexico, west into the highest mountain ranges of the Great Basin. It is composed of barren and sparsely vegetated alpine substrates, typically including both bedrock outcrop and scree slopes, with nonvascular- (lichen) dominated communities. Exposure to desiccating winds, rocky and sometimes unstable substrates, and a short growing season limit plant growth. There can be sparse cover of forbs, grasses, lichens and low shrubs.
S004	<i>Rocky Mountain Alpine Fell-Field</i> - This ecological system is found discontinuously at alpine elevations throughout the Rocky Mountains, west into the mountainous areas of the Great Basin, and north into the Canadian Rockies. Small areas are represented in the westside of the Okanagan Ecoregion in the eastern Cascades. These are wind-scoured fell-fields that are free of snow in the winter, such as ridgetops and exposed saddles, exposing the plants to severe environmental stress. Soils on these windy unproductive sites are shallow, stony, low in organic matter, and poorly developed; wind deflation often results in a gravelly pavement. Most fell-field plants are cushioned, or matted, frequently succulent, flat to the ground in rosettes and often densely haired and thickly cutinized. Plant cover is 15-50%, while exposed rocks make up the rest. Fell-fields are usually within or adjacent to alpine tundra dry meadows. Common species include <i>Arenaria capillaris</i> , <i>Carex albonigra</i> , <i>Carex paysonis</i> , <i>Geum rossii</i> , <i>Kobresia myosuroides</i> , <i>Minuartia obtusiloba</i> , <i>Myosotis asiatica</i> , <i>Paronychia pulvinata</i> , <i>Phlox pulvinata</i> , <i>Sibbaldia procumbens</i> , and <i>Silene acaulis</i> .
S006	<i>Rocky Mountain Cliff and Canyon</i> - This ecological system of barren and sparsely vegetated landscapes (generally <10% plant cover) is found from foothill to subalpine elevations on steep cliff faces, narrow canyons, and smaller rock outcrops of various igneous, sedimentary, and metamorphic bedrock types. It is located throughout the Rocky Mountains and northeastern Cascade Ranges in North America. Also included are unstable scree and talus slopes that typically occur below cliff faces. There may be small patches of dense vegetation, but it typically includes scattered trees and/or shrubs. Characteristic trees includes species from the surrounding landscape, such as <i>Pseudotsuga menziesii</i> , <i>Pinus ponderosa</i> , <i>Pinus flexilis</i> , <i>Populus tremuloides</i> , <i>Abies concolor</i> , <i>Abies lasiocarpa</i> , or <i>Pinus edulis</i> and <i>Juniperus</i> spp. at lower elevations. There may be scattered shrubs present, such as species of <i>Holodiscus</i> , <i>Ribes</i> , <i>Physocarpus</i> , <i>Rosa</i> , <i>Juniperus</i> , and <i>Jamesia americana</i> , <i>Mahonia repens</i> , <i>Rhus trilobata</i> , or <i>Amelanchier alnifolia</i> . Soil development is limited, as is herbaceous cover.
S008	<i>Western Great Plains Cliff and Outcrop</i> - This system includes cliffs and outcrops throughout the Western Great Plains Division. Substrate can range from sandstone and limestone, which can often form bands in the examples of this system. Vegetation is restricted to shelves, cracks and crevices in the rock. However, this system differs from Western Great Plains Badlands (CES303.663) in that often the soil is slightly developed and less erodible, and some grass and shrub species can occur at greater than 10%. Common species in this system include short shrubs such as <i>Rhus trilobata</i> and <i>Artemisia longifolia</i> and mixedgrass species such as <i>Bouteloua curtipendula</i> and <i>Bouteloua gracilis</i> and <i>Calamovilfa longifolia</i> . Drought and wind erosion are the most common natural dynamics affecting this system.
S010	<i>Colorado Plateau Mixed Bedrock Canyon and Tableland</i> - This ecological system is found from foothill to subalpine elevations and includes barren and sparsely vegetated landscapes (generally <10% plant cover) of steep cliff faces, narrow canyons, and smaller rock outcrops of various igneous, sedimentary, and metamorphic bedrock types. Also included is vegetation of unstable scree and talus slopes that typically occurs below cliff faces. Widely scattered trees and shrubs may include <i>Abies concolor</i> , <i>Pinus edulis</i> , <i>Pinus flexilis</i> , <i>Pinus monophylla</i> , <i>Juniperus</i> spp., <i>Artemisia tridentata</i> , <i>Purshia tridentata</i> , <i>Cercocarpus ledifolius</i> , <i>Ephedra</i> spp., <i>Holodiscus discolor</i> , and other species often common in adjacent plant communities.

Appendix D Cont.  
**Code Description**

---

- S011 *Inter-Mountain Basins Shale Badlands* - This widespread ecological system of the intermountain western U.S. is composed of barren and sparsely vegetated substrates (<10% plant cover) typically derived from marine shales but also includes substrates derived from siltstones and mudstones (clay). Landforms are typically rounded hills and plains that form a rolling topography. The harsh soil properties and high rate of erosion and deposition are driving environmental variables supporting sparse dwarf-shrubs, e.g., *Atriplex corrugata*, *Atriplex gardneri*, *Artemisia pedatifida*, and herbaceous vegetation.
- S012 *Inter-Mountain Basins Active and Stabilized Dunes* - This ecological system occurs in Intermountain West basins and is composed of unvegetated to moderately vegetated (<10-30% plant cover) active and stabilized dunes and sandsheets. Species occupying these environments are often adapted to shifting, coarse-textured substrates (usually quartz sand) and form patchy or open grasslands, shrublands or steppe composed of *Achnatherum hymenoides*, *Artemisia filifolia*, *Artemisia tridentata* ssp. *tridentata*, *Atriplex canescens*, *Ephedra* spp., *Coleogyne ramosissima*, *Ericameria nauseosa*, *Leymus flavescens*, *Prunus virginiana*, *Psoralidium lanceolatum*, *Purshia tridentata*, *Sporobolus airoides*, *Tetradymia tetrameres*, or *Tiquilia* spp.
- S013 *Inter-Mountain Basins Volcanic Rock and Cinder Land* - This ecological system occurs in the intermountain western U.S. and is limited to barren and sparsely vegetated volcanic substrates (generally <10% plant cover) such as basalt lava (malpais), basalt dikes with associated colluvium, basalt cliff faces and uplifted "backbones," tuff, cinder cones or cinder fields. It may occur as large-patch, small-patch and linear (dikes) spatial patterns. Vegetation is variable and includes a variety of species depending on local environmental conditions, e.g., elevation, age and type of substrate. At montane and foothill elevations scattered *Pinus ponderosa*, *Pinus flexilis*, or *Juniperus* spp. trees may be present. Shrubs such as *Ephedra* spp., *Atriplex canescens*, *Eriogonum corymbosum*, *Eriogonum ovalifolium*, and *Fallugia paradoxa* are often present on some lava flows and cinder fields. Species typical of sand dunes such as *Andropogon hallii* and *Artemisia filifolia* may be present on cinder substrates.
- S014<sup>4</sup> *Inter-Mountain Basins Greasewood Wash* - This barren and sparsely vegetated (generally <10% plant cover) ecological system is restricted to intermittently flooded streambeds and banks that are often lined with shrubs such as *Sarcobatus vermiculatus*, *Ericameria nauseosa*, *Fallugia paradoxa*, and/or *Artemisia cana* ssp. *cana* (in more northern and mesic stands). *Grayia spinosa* may dominate in the Great Basin. Shrubs form a continuous or intermittent linear canopy in and along drainages but do not extend out into flats. Typically it includes patches of saltgrass meadow where water remains for the longest periods. Soils are generally less alkaline than those found in the playa system. Desert scrub species (e.g., *Acacia greggii*, *Prosopis* spp.), that are common in the Mojave, Sonoran and Chihuahuan desert washes, are not present. This type can occur in limited portions of the southwestern Great Plains.
- S015<sup>4</sup> *Inter-Mountain Basins Playa* - This ecological system is composed of barren and sparsely vegetated playas (generally <10% plant cover) found in the intermountain western U.S. Salt crusts are common throughout, with small saltgrass beds in depressions and sparse shrubs around the margins. These systems are intermittently flooded. The water is prevented from percolating through the soil by an impermeable soil subhorizon and is left to evaporate. Soil salinity varies greatly with soil moisture and greatly affects species composition. Characteristic species may include *Allenrolfea occidentalis*, *Sarcobatus vermiculatus*, *Grayia spinosa*, *Puccinellia lemmonii*, *Leymus cinereus*, *Distichlis spicata*, and/or *Atriplex* spp.
- S016 *North American Warm Desert Bedrock Cliff and Outcrop* - This ecological system is found from subalpine to foothill elevations and includes barren and sparsely vegetated landscapes (generally <10% plant cover) of steep cliff faces, narrow canyons, and smaller rock outcrops of various igneous, sedimentary, and metamorphic bedrock types. Also included are unstable scree and talus slopes that typically occur bellow cliff faces. Species present are diverse and may include *Bursera microphylla*, *Fouquieria splendens*, *Nolina bigelovii*, *Opuntia bigelovii*, and other desert species, especially succulents. Lichens are predominant lifeforms in some areas. May include a variety of desert shrublands less than 2 ha (5 acres) in size from adjacent areas.

Appendix D Cont.  
Code Description

- S018 *North American Warm Desert Active and Stabilized Dunes* - This ecological system occurs across the warm deserts of North America and is composed of unvegetated to sparsely vegetated (generally <10% plant cover) active dunes and sandsheets derived from quartz or gypsum sands. Common vegetation includes *Ambrosia dumosa*, *Abronia villosa*, *Eriogonum deserticola*, *Larrea tridentata*, *Pleuraphis rigida*, *Poliomintha* spp., *Prosopis* spp., *Psoralea* spp., *Artemisia filifolia*, and *Rhus microphylla*. Dune "blowouts" and subsequent stabilization through succession are characteristic processes.
- S019 *North American Warm Desert Volcanic Rockland* - This ecological system occurs across the warm deserts of North America and is restricted to barren and sparsely vegetated (<10% plant cover) volcanic substrates such as basalt lava (malpais) and tuff. Vegetation is variable and includes a variety of species depending on local environmental conditions, e.g., elevation, age and type of substrate. Typically scattered *Larrea tridentata*, *Atriplex hymenelytra*, or other desert shrubs are present.
- S020<sup>4</sup> *North American Warm Desert Wash* - This ecological system is restricted to intermittently flooded washes or arroyos that dissect bajadas, mesas, plains and basin floors throughout the warm deserts of North America. Although often dry, the intermittent fluvial processes define this system, which are often associated with rapid sheet and gully flow. This system occurs as linear or braided strips within desert scrub- or desert grassland-dominated landscapes. The vegetation of desert washes is quite variable ranging from sparse and patchy to moderately dense and typically occurs along the banks, but may occur within the channel. The woody layer is typically intermittent to open and may be dominated by shrubs and small trees such as *Acacia greggii*, *Brickellia laciniata*, *Baccharis sarothroides*, *Chilopsis linearis*, *Fallugia paradoxa*, *Hymenoclea salsola*, *Hymenoclea monogyra*, *Juglans microcarpa*, *Prosopis* spp., *Psoralea spinosus*, *Prunus fasciculata*, *Rhus microphylla*, *Salazaria mexicana*, or *Sarcobatus vermiculatus*.
- S021 *North American Warm Desert Pavement* - This ecological system occurs throughout much of the warm deserts of North America and is composed of unvegetated to very sparsely vegetated (<2% plant cover) landscapes, typically flat basins where extreme temperature and wind develop ground surfaces of fine to medium gravel coated with "desert varnish." Very low cover of desert scrub species such as *Larrea tridentata* or *Eriogonum fasciculatum* is usually present. However, ephemeral herbaceous species may have high cover in response to seasonal precipitation, including *Chorizanthe rigida*, *Eriogonum inflatum*, and *Geraea canescens*.
- S022<sup>4</sup> *North American Warm Desert Playa* - This system is composed of barren and sparsely vegetated playas (generally <10% plant cover) found across the warm deserts of North America, extending into the extreme southern end of the San Joaquin Valley in California. Playas form with intermittent flooding, followed by evaporation, leaving behind a saline residue. Salt crusts are common throughout, with small saltgrass beds in depressions and sparse shrubs around the margins. Subsoils often include an impermeable layer of clay or caliche. Large desert playas tend to be defined by vegetation rings formed in response to salinity. Given their common location in wind-swept desert basins, dune fields often form downwind of large playas. In turn, playas associated with dunes often have a deeper water supply. Species may include *Allenrolfea occidentalis*, *Suaeda* spp., *Distichlis spicata*, *Eleocharis palustris*, *Oryzopsis* spp., *Sporobolus* spp., *Tiquilia* spp., or *Atriplex* spp. Ephemeral herbaceous species may have high cover periodically. Adjacent vegetation is typically Sonora-Mojave Desert Mixed Salt Desert Scrub (CES302.749), Chihuahuan Mixed Salt Desert Scrub (CES302.017), Gulf of California Coastal Mixed Salt Desert Scrub (CES302.015), Baja California del Norte Gulf Coast Ocotillo-Limberbush-Creosotebush Desert Scrub (CES302.014), or Chihuahuan Creosotebush Basin Desert Scrub (CES302.731).
- S023 *Rocky Mountain Aspen Forest and Woodland* - This widespread ecological system is more common in the southern and central Rocky Mountains, but occurs throughout much of the western U.S. and north into Canada, in the montane and subalpine zones. Elevations generally range from 1525 to 3050 m (5000-10,000 feet), but occurrences can be found at lower elevations in some regions. Distribution of this ecological system is primarily limited by adequate soil moisture required to meet its high evapotranspiration demand, and secondarily is limited by the length of the growing season or low temperatures. These are upland forests

Appendix D Cont.  
**Code Description**

---

- S023 and woodlands dominated by *Populus tremuloides* without a significant conifer component (<25% relative tree cover). The understory structure may be complex with multiple shrub and herbaceous layers, or simple with just an herbaceous layer. The herbaceous layer may be dense or sparse, dominated by graminoids or forbs. Associated shrub species include *Symphoricarpos* spp., *Rubus parviflorus*, *Amelanchier alnifolia*, and *Arctostaphylos uva-ursi*. Occurrences of this system originate and are maintained by stand-replacing disturbances such as avalanches, crown fire, insect outbreak, disease and windthrow, or clearcutting by man or beaver, within the matrix of conifer forests.
- S024<sup>4</sup> *Rocky Mountain Bigtooth Maple Ravine Woodland* - This ecological system occurs in cool ravines, on toeslopes and slump benches associated with riparian areas in the northern and central Wasatch Range and Tavaputs Plateau extending into southern Idaho, as well as in scattered localities in southwestern Utah, central Arizona and New Mexico and the Trans-Pecos of Texas. Substrates are typically rocky colluvial or alluvial soils with favorable soil moisture. These woodlands are dominated by *Acer grandidentatum* but may include mixed stands codominated by *Quercus gambelii* or with scattered conifers. Some stands may include *Acer negundo* or *Populus tremuloides* as minor components. It also occurs on steeper, north-facing slopes at higher elevations, often adjacent to Rocky Mountain Gambel Oak-Mixed Montane Shrubland (CES306.818) or Rocky Mountain Aspen Forest and Woodland (CES306.813).
- S025 *Rocky Mountain Subalpine-Montane Limber-Bristlecone Pine Woodland* - This ecological system occurs throughout the Rocky Mountains on dry, rocky ridges and slopes near upper treeline above the matrix spruce-fir forest. It extends down to the lower montane in the central and northern Rocky Mountains and northeastern Great Basin mountains where dominated by *Pinus flexilis*, particularly along the Front Range north into Canada. Sites are harsh, exposed to desiccating winds, with rocky substrates and a short growing season that limit plant growth. Higher-elevation occurrences are found well into the subalpine-alpine transition on wind-blasted, mostly westfacing slopes and exposed ridges. Calcareous substrates are important for *Pinus flexilis*-dominated communities in the northern Rocky Mountains and possibly elsewhere. The open tree canopy is often patchy and is strongly dominated by *Pinus flexilis* or *Pinus aristata* with the latter restricted to southern Colorado, northern New Mexico and the San Francisco Mountains in Arizona. In the northern Rockies and northern Great Basin, *Pinus albicaulis* is found in some occurrences. Other trees such as *Juniperus* spp., *Pinus contorta*, *Pinus ponderosa*, or *Pseudotsuga menziesii* are occasionally present. *Arctostaphylos uva-ursi*, *Cercocarpus ledifolius*, *Juniperus communis*, *Mahonia repens*, *Purshia tridentata*, *Ribes montigenum*, or *Vaccinium* spp. may form an open shrub layer in some stands. The herbaceous layer, if present, is generally sparse and composed of xeric graminoids, such as *Calamagrostis purpurascens*, *Festuca arizonica*, *Festuca idahoensis*, *Festuca thurberi*, or *Pseudoroegneria spicata*, or more alpine plants.
- S028 *Rocky Mountain Subalpine Dry-Mesic Spruce-Fir Forest and Woodland* - Engelmann spruce and subalpine fir forests comprise a substantial part of the subalpine forests of the Cascades and Rocky Mountains from southern British Columbia east into Alberta, south into New Mexico and the Intermountain region. They are the matrix forests of the subalpine zone, with elevations ranging from 1275 m in its northern distribution to 3355 m in the south (4100-11,000 feet). They often represent the highest elevation forests in an area. Sites within this system are cold year-round, and precipitation is predominantly in the form of snow, which may persist until late summer. Snowpacks are deep and late-lying, and summers are cool. Frost is possible almost all summer and may be common in restricted topographic basins and benches. Despite their wide distribution, the tree canopy characteristics are remarkably similar, with *Picea engelmannii* and *Abies lasiocarpa* dominating either mixed or alone. *Pseudotsuga menziesii* may persist in occurrences of this system for long periods without regeneration. *Pinus contorta* is common in many occurrences, and patches of pure *Pinus contorta* are not uncommon, as well as mixed conifer/*Populus tremuloides* stands. In some areas, such as Wyoming, *Picea engelmannii*-dominated forests are on limestone or dolomite, while nearby codominated spruce-fir forests are on granitic or volcanic rocks. Xeric species may include *Juniperus communis*, *Linnaea borealis*, *Mahonia repens*, or *Vaccinium scoparium*. More northern occurrences often have taller, more mesic shrub and herbaceous species, such as *Empetrum nigrum*, *Rhododendron albiflorum*, and *Vaccinium membranaceum*. Disturbance includes occasional blow-down, insect outbreaks and stand-replacing fire.

Appendix D Cont.  
Code Description

- S030 *Rocky Mountains Subalpine Mesic Spruce-fir Forest and Woodland* - This is a high-elevation system of the Rocky Mountains, dominated by *Picea engelmannii* and *Abies lasiocarpa*. It extends eastward into the northeastern Olympic Mountains and the northeastern side of Mount Rainier in Washington. Occurrences are typically found in locations with cold-air drainage or ponding, or where snowpacks linger late into the summer, such as north-facing slopes and high-elevation ravines. They can extend down in elevation below the subalpine zone in places where cold-air ponding occurs; northerly and easterly aspects predominate. These forests are found on gentle to very steep mountain slopes, high-elevation ridgetops and upper slopes, plateau-like surfaces, basins, alluvial terraces, well-drained benches, and inactive stream terraces. In the Olympics and northern Cascades, the climate is more maritime than typical for this system, but due to the lower snowfall in these rainshadow areas, summer drought may be more significant than snowpack in limiting tree regeneration in burned areas. *Picea engelmannii* is rare in these areas. Mesic understory shrubs include *Menziesia ferruginea*, *Vaccinium membranaceum*, *Rhododendron albiflorum*, *Amelanchier alnifolia*, *Rubus parviflorus*, *Ledum glandulosum*, *Phyllodoce empetriformis*, and *Salix* spp. Herbaceous species include *Actaea rubra*, *Maianthemum stellatum*, *Cornus canadensis*, *Erigeron eximius*, *Gymnocarpium dryopteris*, *Rubus pedatus*, *Saxifraga bronchialis*, *Tiarella* spp., *Lupinus arcticus* ssp. *subalpinus*, *Valeriana sitchensis*, and graminoids *Luzula glabrata* var. *hitchcockii* or *Calamagrostis canadensis*. Disturbances include occasional blow-down, insect outbreaks and stand-replacing fire.
- S031 *Rocky Mountains Lodgepole Pine Forest* - This system is widespread in upper montane to subalpine elevations of the Rocky Mountains, Intermountain region, and north into the Canadian Rockies. These are subalpine forests where the dominance of *Pinus contorta* is related to fire history and topo-edaphic conditions. Following stand-replacing fires, *Pinus contorta* will rapidly colonize and develop into dense, even-aged stands. Most forests in this ecological system are early- to mid-successional forests which developed following fires. Some *Pinus contorta* forests will persist on sites that are too extreme for other conifers to establish. These include excessively well-drained pumice deposits, glacial till and alluvium on valley floors where there is cold air accumulation, warm and droughty shallow soils over fractured quartzite bedrock, and shallow moisture-deficient soils with a significant component of volcanic ash. Soils supporting these forests are typically well-drained, gravelly, coarse-textured, acidic, and rarely formed from calcareous parent materials. These forests are dominated by *Pinus contorta* with shrub, grass, or barren understories. Sometimes there are intermingled mixed conifer/*Populus tremuloides* stands with the latter occurring with inclusions of deeper, typically fine-textured soils. The shrub stratum may be conspicuous to absent; common species include *Arctostaphylos uva-ursi*, *Ceanothus velutinus*, *Linnaea borealis*, *Mahonia repens*, *Purshia tridentata*, *Spiraea betulifolia*, *Spiraea douglasii*, *Shepherdia canadensis*, *Vaccinium caespitosum*, *Vaccinium scoparium*, *Vaccinium membranaceum*, *Symphoricarpos albus*, and *Ribes* spp. In southern interior British Columbia, this system is usually an open lodgepole pine forest found extensively between 500 and 1600 m elevation in the Columbia range. In the Interior Cedar Hemlock and Interior Douglas-fir zones, *Tsuga heterophylla* or *Pseudotsuga menziesii* may be present.
- S032<sup>5</sup> *Rocky Mountain Montane Dry-Mesic Mixed Conifer Forest and Woodland* - This is a highly variable ecological system of the montane zone of the Rocky Mountains. It occurs throughout the southern Rockies, north and west into Utah, Nevada, western Wyoming and Idaho. These are mixed-conifer forests occurring on all aspects at elevations ranging from 1200 to 3300 m. Rainfall averages less than 75 cm per year (40-60 cm) with summer "monsoons" during the growing season contributing substantial moisture. The composition and structure of overstory is dependent upon the temperature and moisture relationships of the site, and the successional status of the occurrence. *Pseudotsuga menziesii* and *Abies concolor* are most frequent, but *Pinus ponderosa* may be present to codominant. *Pinus flexilis* is common in Nevada. *Pseudotsuga menziesii* forests occupy drier sites, and *Pinus ponderosa* is a common codominant. *Abies concolor*-dominated forests occupy cooler sites, such as upper slopes at higher elevations, canyon sideslopes, ridgetops, and north- and east-facing slopes which burn somewhat infrequently. *Picea pungens* is most often found in cool, moist locations, often occurring as smaller patches within a matrix of other associations. As many as seven conifers can be found growing in the same occurrence, and there are a number of cold-deciduous shrub and graminoid species common, including *Arctostaphylos uva-ursi*, *Mahonia repens*, *Paxistima myrsinites*, *Symphoricarpos oreophilus*, *Jamesia americana*, *Quercus gambelii*, and *Festuca arizonica*. This system was

Appendix D Cont.  
**Code Description**

---

- S032<sup>5</sup> undoubtedly characterized by a mixed severity fire regime in its "natural condition," characterized by a high degree of variability in lethality and return interval.
- S034<sup>5</sup> *Rocky Mountain Montane Mesic Mixed Conifer Forest and Woodland* - These are mixed-conifer forests of the Rocky Mountains west into the ranges of the Great Basin, occurring predominantly in cool ravines and on north-facing slopes. Elevations range from 1200 to 3300 m. Occurrences of this system are found on cooler and more mesic sites than Rocky Mountain Montane Dry-Mesic Mixed Conifer Forest and Woodland (CES306.823). Such sites include lower and middle slopes of ravines, along stream terraces, moist, concave topographic positions and north- and east-facing slopes which burn somewhat infrequently. *Pseudotsuga menziesii* and *Abies concolor* are most common canopy dominants, but *Picea engelmannii*, *Picea pungens*, or *Pinus ponderosa* may be present. This system includes mixed conifer/*Populus tremuloides* stands. A number of cold-deciduous shrub species can occur, including *Acer glabrum*, *Acer grandidentatum*, *Alnus incana*, *Betula occidentalis*, *Cornus sericea*, *Jamesia americana*, *Physocarpus malvaceus*, *Robinia neomexicana*, *Vaccinium membranaceum*, and *Vaccinium myrtillus*. Herbaceous species include *Bromus ciliatus*, *Carex geyeri*, *Carex rossii*, *Carex siccata*, *Muhlenbergia virescens*, *Pseudoroegneria spicata*, *Erigeron eximius*, *Fragaria virginiana*, *Luzula parviflora*, *Osmorhiza berteroi*, *Packera cardamine*, *Thalictrum occidentale*, and *Thalictrum fendleri*. Naturally occurring fires are of variable return intervals, and mostly light, erratic, and infrequent due to the cool, moist conditions.
- S035<sup>3</sup> *Madrean Pine-Oak Forest and Woodland* - This system occurs on mountains and plateaus in the Sierra Madre Occidentale and Sierra Madre Orientale in Mexico, Trans-Pecos Texas, southern New Mexico and Arizona, generally south of the Mogollon Rim. These forests and woodlands are composed of Madrean pines (*Pinus arizonica*, *Pinus engelmannii*, *Pinus leiophylla*, or *Pinus strobiformis*) and evergreen oaks (*Quercus arizonica*, *Quercus emoryi*, or *Quercus grisea*) intermingled with patchy shrublands on most mid-elevation slopes (1500-2300 m elevation). Other tree species include *Cupressus arizonica*, *Juniperus deppeana*, *Pinus cembroides*, *Pinus discolor*, *Pinus ponderosa* (with Madrean pines or oaks), and *Pseudotsuga menziesii*. Subcanopy and shrub layers may include typical encinal and chaparral species such as *Agave* spp., *Arbutus arizonica*, *Arctostaphylos pringlei*, *Arctostaphylos pungens*, *Garrya wrightii*, *Nolina* spp., *Quercus hypoleucoides*, *Quercus rugosa*, and *Quercus turbinella*. Some stands have moderate cover of perennial graminoids such as *Muhlenbergia emersleyi*, *Muhlenbergia longiligula*, *Muhlenbergia virescens*, and *Schizachyrium cirratum*. Fires are frequent with perhaps more crown fires than ponderosa pine woodlands, which tend to have more frequent ground fires on gentle slopes.
- S036 *Rocky Mountain Ponderosa Pine Woodland* - This very widespread ecological system is most common throughout the cordillera of the Rocky Mountains, from the Greater Yellowstone region south. It is also found in the Colorado Plateau region, west into scattered locations in the Great Basin, and in the Black Hills of South Dakota and Wyoming. These woodlands occur at the lower treeline/ecotone between grassland or shrubland and more mesic coniferous forests typically in warm, dry, exposed sites. Elevations range from less than 1900 m in northern Wyoming to 2800 m in the New Mexico mountains. Occurrences are found on all slopes and aspects; however, moderately steep to very steep slopes or ridgetops are most common. This ecological system generally occurs on igneous, metamorphic, and sedimentary material derived soils, with characteristic features of good aeration and drainage, coarse textures, circumneutral to slightly acid pH, an abundance of mineral material, rockiness, and periods of drought during the growing season. Northern Rocky Mountain Ponderosa Pine Woodland (CES306.030) in the eastern Cascades, Okanagan and northern Rockies regions receives winter and spring rains, and thus has a greater spring "green-up" than the drier woodlands in the central Rockies. *Pinus ponderosa* (primarily var. *scopulorum* and var. *brachyptera*) is the predominant conifer; *Pseudotsuga menziesii*, *Pinus edulis*, and *Juniperus* spp. may be present in the tree canopy. The understory is usually shrubby, with *Artemisia nova*, *Artemisia tridentata*, *Arctostaphylos patula*, *Arctostaphylos uva-ursi*, *Cercocarpus montanus*, *Purshia stansburiana*, *Purshia tridentata*, *Quercus gambelii*, *Symphoricarpos oreophilus*, *Prunus virginiana*, *Amelanchier alnifolia*, and *Rosa* spp. common species. *Pseudoroegneria spicata* and species of *Hesperostipa*, *Achnatherum*, *Festuca*, *Muhlenbergia*, and *Bouteloua* are some of the common grasses. Mixed fire regimes and ground fires of variable return intervals maintain these woodlands, depending on climate, degree of soil development, and understory density.

Appendix D Cont.  
Code Description

- S038 *Southern Rocky Mountain Pinyon-Juniper Woodland* - This southern Rocky Mountain ecological system occurs on dry mountains and foothills in southern Colorado east of the Continental Divide, in mountains and plateaus of north-central New Mexico, and extends out onto limestone breaks in the southeastern Great Plains. These woodlands occur on warm, dry sites on mountain slopes, mesas, plateaus, and ridges. Severe climatic events occurring during the growing season, such as frosts and drought, are thought to limit the distribution of pinyon-juniper woodlands to relatively narrow altitudinal belts on mountainsides. Soils supporting this system vary in texture ranging from stony, cobbly, gravelly sandy loams to clay loam or clay. *Pinus edulis* and/or *Juniperus monosperma* dominate the tree canopy. *Juniperus scopulorum* may codominate or replace *Juniperus monosperma* at higher elevations. Stands with *Juniperus osteosperma* are representative the Colorado Plateau and are not included in this system. In southern transitional areas between Madrean Pinyon-Juniper Woodland (CES305.797) and Southern Rocky Mountain Pinyon-Juniper Woodland (CES306.835) in central New Mexico, *Juniperus deppeana* becomes common. Understory layers are variable and may be dominated by shrubs, graminoids, or be absent. Associated species are more typical of southern Rocky Mountains than the Colorado Plateau and include *Artemisia bigelovii*, *Cercocarpus montanus*, *Quercus gambelii*, *Achnatherum scribneri*, *Bouteloua gracilis*, *Festuca arizonica*, or *Pleuraphis jamesii*.
- S039 *Colorado Plateau Pinyon-Juniper Woodland* - This ecological system occurs in dry mountains and foothills of the Colorado Plateau region including the Western Slope of Colorado to the Wasatch Range, south to the Mogollon Rim and east into the northwestern corner of New Mexico. It is typically found at lower elevations ranging from 1500-2440 m. These woodlands occur on warm, dry sites on mountain slopes, mesas, plateaus, and ridges. Severe climatic events occurring during the growing season, such as frosts and drought, are thought to limit the distribution of pinyon-juniper woodlands to relatively narrow altitudinal belts on mountainsides. Soils supporting this system vary in texture ranging from stony, cobbly, gravelly sandy loams to clay loam or clay. *Pinus edulis* and/or *Juniperus osteosperma* dominate the tree canopy. In the southern portion of the Colorado Plateau in northern Arizona and northwestern New Mexico, *Juniperus monosperma* and hybrids of *Juniperus* spp may dominate or codominate the tree canopy. *Juniperus scopulorum* may codominate or replace *Juniperus osteosperma* at higher elevations. Understory layers are variable and may be dominated by shrubs, graminoids, or be absent. Associated species include *Arctostaphylos patula*, *Artemisia tridentata*, *Cercocarpus intricatus*, *Cercocarpus montanus*, *Coleogyne ramosissima*, *Purshia stansburiana*, *Purshia tridentata*, *Quercus gambelii*, *Bouteloua gracilis*, *Pleuraphis jamesii*, or *Poa fendleriana*. This system occurs at higher elevations than Great Basin Pinyon-Juniper Woodland (CES304.773) and Colorado Plateau shrubland systems where sympatric.
- S042 *Inter-Mountain Basins Aspen-Mixed Conifer Forest and Woodland* - This ecological system occurs on montane slopes and plateaus in Utah, western Colorado, northern Arizona, eastern Nevada, southern Idaho and western Wyoming. Elevations range from 1700 to 2800 m. Occurrences are typically on gentle to steep slopes on any aspect but are often found on clay-rich soils in intermontane valleys. Soils are derived from alluvium, colluvium and residuum from a variety of parent materials but most typically occur on sedimentary rocks. The tree canopy is composed of a mix of deciduous and coniferous species, codominated by *Populus tremuloides* and conifers, including *Pseudotsuga menziesii*, *Abies concolor*, *Abies lasiocarpa*, *Picea engelmannii*, *Picea pungens*, *Pinus contorta*, *Pinus flexilis*, and *Pinus ponderosa*. As the occurrences age, *Populus tremuloides* is slowly reduced until the conifer species become dominant. Common shrubs include *Amelanchier alnifolia*, *Prunus virginiana*, *Acer grandidentatum*, *Symphoricarpos oreophilus*, *Juniperus communis*, *Paxistima myrsinites*, *Rosa woodsii*, *Spiraea betulifolia*, *Symphoricarpos albus*, or *Mahonia repens*. Herbaceous species include *Bromus carinatus*, *Calamagrostis rubescens*, *Carex geyeri*, *Elymus glaucus*, *Poa* spp., and *Achnatherum*, *Hesperostipa*, *Nassella*, and/or *Piptochaetium* spp. (= *Stipa* spp.), *Achillea millefolium*, *Arnica cordifolia*, *Asteraceae* spp., *Erigeron* spp., *Galium boreale*, *Geranium viscosissimum*, *Lathyrus* spp., *Lupinus argenteus*, *Mertensia arizonica*, *Mertensia lanceolata*, *Maianthemum stellatum*, *Osmorhiza berteroi* (= *Osmorhiza chilensis*), and *Thalictrum fendleri*. Most occurrences at present represent a late-seral stage of aspen changing to a pure conifer occurrence. Nearly a hundred years of fire suppression and livestock grazing have converted much of the pure aspen occurrences to the present-day aspen-conifer forest and woodland ecological system.

Appendix D Cont.  
**Code Description**

---

- S042 cont. In order to capture important habitat characteristics of an aspen-mixed conifer ecological system for vertebrate habitat modeling, SWReGAP land cover mappers mapped patches of aspen-mixed conifer stands outside its normal range into the Southern Rocky Mountains. In the Southern Rocky Mountains, this system occurs as small to large patches of aspenmixed conifer woodland that could also be interpreted as seral stands within several Rocky Mountain conifer forest and woodland systems including: S028 Rocky Mountain Subalpine Dry-Mesic Spruce-Fir Forest and Woodland, S030 Rocky Mountain Subalpine Mesic Spruce-Fir Forest and Woodland, S031 Rocky Mountain Lodgepole Pine Forest, S032 Rocky Mountain Montane Dry-Mesic Mixed Conifer Forest and Woodland, S034 Rocky Mountain Montane Mesic Mixed Conifer Forest and Woodland, and S036 Rocky Mountain Ponderosa Pine Woodland (see individual descriptions for additional information).
- S046 *Rocky Mountain Gambel Oak-Mixed Montane Shrubland* - This ecological system occurs in the mountains, plateaus and foothills in the southern Rocky Mountains and Colorado Plateau including the Uinta and Wasatch ranges and the Mogollon Rim. These shrublands are most commonly found along dry foothills, lower mountain slopes, and at the edge of the western Great Plains from approximately 2000 to 2900 m in elevation, and are often situated above pinyon-juniper woodlands. Substrates are variable and include soil types ranging from calcareous, heavy, fine-grained loams to sandy loams, gravelly loams, clay loams, deep alluvial sand, or coarse gravel. The vegetation is typically dominated by *Quercus gambelii* alone or codominant with *Amelanchier alnifolia*, *Amelanchier utahensis*, *Artemisia tridentata*, *Cercocarpus montanus*, *Prunus virginiana*, *Purshia stansburiana*, *Purshia tridentata*, *Robinia neomexicana*, *Symphoricarpos oreophilus*, or *Symphoricarpos rotundifolius*. There may be inclusions of other mesic montane shrublands with *Quercus gambelii* absent or as a relatively minor component. This ecological system intergrades with the lower montane-foothills shrubland system and shares many of the same site characteristics. Density and cover of *Quercus gambelii* and *Amelanchier* spp. often increase after fire.
- S047 *Rocky Mountains Lower Montane-Foothill Shrubland* - This ecological system is found in the foothills, canyon slopes and lower mountains of the Rocky Mountains and on outcrops and canyon slopes in the western Great Plains. It ranges from southern New Mexico extending north into Wyoming, and west into the Intermountain region. These shrublands occur between 1500-2900 m elevations and are usually associated with exposed sites, rocky substrates, and dry conditions, which limit tree growth. It is common where *Quercus gambelii* is absent such as the northern Colorado Front Range and in drier foothills and prairie hills. This system is generally drier than Rocky Mountain Gambel Oak-Mixed Montane Shrubland (CES306.818), but may include mesic montane shrublands where *Quercus gambelii* does not occur. Scattered trees or inclusions of grassland patches or steppe may be present, but the vegetation is typically dominated by a variety of shrubs including *Amelanchier utahensis*, *Cercocarpus montanus*, *Purshia tridentata*, *Rhus trilobata*, *Ribes cereum*, *Symphoricarpos oreophilus*, or *Yucca glauca*. In northeastern Wyoming and north into adjacent Montana, *Cercocarpus ledifolius*, usually with *Artemisia tridentata*, is the common dominant shrub. Grasses are represented as species of *Muhlenbergia*, *Bouteloua*, *Hesperostipa*, and *Pseudoroegneria spicata*. Fires play an important role in this system as the dominant shrubs usually have a severe dieback, although some plants will stump sprout. *Cercocarpus montanus* requires a disturbance such as fire to reproduce, either by seed sprout or root crown sprouting. Fire suppression may have allowed an invasion of trees into some of these shrublands, but in many cases sites are too xeric for tree growth.
- S048<sup>1</sup> *Western Great Plains Sandhill Sagebrush Shrubland* - This system is found mostly in south-central areas of the Western Great Plains Division ranging from the Nebraska Sandhill region south to central Texas, although some examples may reach as far north as the Badlands of South Dakota. The climate is semi-arid to arid for much of the region in which this system occurs. This system is found on somewhat excessively to excessively well-drained, deep sandy soils that are often associated with dune systems and ancient floodplains. In some areas, this system may actually occur as a result of overgrazing in Western Great Plains Tallgrass Prairie (CES303.673) or Western Great Plains Sand Prairie (CES303.670). This system is characterized by a sparse to moderately dense woody layer dominated by *Artemisia filifolia*. Associated species can vary with geography, amount and season of precipitation, disturbance and soil texture. Several graminoid species such as *Andropogon hallii*, *Schizachyrium scoparium*, *Sporobolus cryptandrus*,

Appendix D Cont.  
Code Description

- S048<sup>1</sup> Calamovilfa gigantea, Hesperostipa comata, and Bouteloua spp. can be connected with this system. Other shrub species may also be present including Yucca glauca, Prosopis glandulosa, Rhus trilobata, and Prunus angustifolia. In the southern range of this system, Quercus havardii may also be present and represents one succession pathway that develops over time following a disturbance. Quercus havardii is able to resprout following a fire and thus may persist for long periods of time once established. Fire and grazing are the most important dynamic processes for this type, although drought stress can impact this system significantly in some areas. Overgrazing can lead to decreasing dominance of some of the grass species such as Andropogon hallii, Calamovilfa gigantea, and Schizachyrium scoparium.
- S051<sup>1</sup> *Madrean Encinal* - Madrean Encinal occurs on foothills, canyons, bajadas and plateaus in the Sierra Madre Occidentale and Sierra Madre Orientale in Mexico, extending north into Trans-Pecos Texas, southern New Mexico and sub-Mogollon Arizona. These woodlands are dominated by Madrean evergreen oaks along a low-slope transition below Madrean Pine-Oak Forest and Woodland (CES305.796) and Madrean Pinyon-Juniper Woodland (CES305.797). Lower elevation stands are typically open woodlands or savannas where they transition into desert grasslands, chaparral or in some cases desertscrub. Common evergreen oak species include Quercus arizonica, Quercus emoryi, Quercus intricata, Quercus grisea, Quercus oblongifolia, Quercus toumeyi, and in Mexico Quercus chihuahuensis and Quercus albocincta. Madrean pine, Arizona cypress, pinyon and juniper trees may be present, but do not codominate. Chaparral species such as Arcostaphylos pungens, Cercocarpus montanus, Purshia spp., Garrya wrightii, Quercus turbinella, Frangula betulifolia (= Rhamnus betulifolia), or Rhus spp. may be present but do not dominate. The graminoid layer is usually prominent between trees in grassland or steppe that is dominated by warm-season grasses such as Aristida spp., Bouteloua gracilis, Bouteloua curtipendula, Bouteloua rothrockii, Digitaria californica, Eragrostis intermedia, Hilaria belangeri, Leptochloa dubia, Muhlenbergia spp., Pleuraphis jamesii, or Schizachyrium cirratum, species typical of Chihuahuan Piedmont Semi-Desert Grassland (CES302.735). This system includes seral stands dominated by shrubby Madrean oaks typically with a strong graminoid layer. In transition areas with drier chaparral systems, stands of chaparral are not dominated by Madrean oaks; however, Madrean Encinal may extend down along drainages.
- S054<sup>1</sup> *Inter-Mountain Basins Big Sagebrush Shrubland* - This ecological system occurs throughout much of the western U.S., typically in broad basins between mountain ranges, plains and foothills between 1500 and 2300 m elevation. Soils are typically deep, well-drained and non-saline. These shrublands are dominated by Artemisia tridentata ssp. tridentata and/or Artemisia tridentata ssp. wyomingensis. Scattered Juniperus spp., Sarcobatus vermiculatus, and Atriplex spp. may be present in some stands. Ericameria nauseosa, Chrysothamnus viscidiflorus, Purshia tridentata, or Symphoricarpos oreophilus may codominate disturbed stands. Perennial herbaceous components typically contribute less than 25% vegetative cover. Common graminoid species include Achnatherum hymenoides, Bouteloua gracilis, Elymus lanceolatus, Festuca idahoensis, Hesperostipa comata, Leymus cinereus, Pleuraphis jamesii, Pascopyrum smithii, Poa secunda, or Pseudoroegneria spicata.
- S056 *Colorado Plateau Mixed Low Sagebrush Shrubland* - This ecological system occurs in the Colorado Plateau, Tavaputs Plateau and Uinta Basin in canyons, gravelly draws, hilltops, and dry flats at elevations generally below 1800 m. Soils are often rocky, shallow, and alkaline. This type extends across northern New Mexico into the southern Great Plains on limestone hills. It includes open shrublands and steppe dominated by Artemisia nova or Artemisia bigelovii sometimes with Artemisia tridentata ssp. wyomingensis codominant. Semi-arid grasses such as Achnatherum hymenoides, Aristida purpurea, Bouteloua gracilis, Hesperostipa comata, Pleuraphis jamesii, or Poa fendleriana are often present and may form a graminoid layer with over 25% cover.
- S057 *Mogollon Chaparral* - This ecological system occurs across central Arizona (Mogollon Rim), western New Mexico and southern Utah and Nevada. It often dominates along the mid-elevation transition from the Mojave, Sonoran, and northern Chihuahuan deserts into mountains (1000-2200 m). It occurs on foothills, mountain slopes and canyons in drier habitats below the encinal and Pinus ponderosa woodlands. Stands are often associated with more xeric and coarse-textured substrates such as limestone, basalt or alluvium,

Appendix D Cont.  
**Code Description**

---

- S057 especially in transition areas with more mesic woodlands. The moderate to dense shrub canopy includes species such as *Quercus turbinella*, *Quercus toumeyii*, *Cercocarpus montanus*, *Canotia holacantha*, *Ceanothus greggii*, *Forestiera pubescens* (= *Forestiera neomexicana*), *Garrya wrightii*, *Juniperus deppeana*, *Purshia stansburiana*, *Rhus ovata*, *Rhus trilobata*, and *Arctostaphylos pungens* and *Arctostaphylos pringlei* at higher elevations. Most chaparral species are fire-adapted, resprouting vigorously after burning or producing fire-resistant seeds. Stands occurring within montane woodlands are seral and a result of recent fires.
- S058 *Chihuahuan Mesquite Upland Scrub* - This ecological system occurs as upland shrublands that are concentrated in the extensive grassland-shrubland transition in foothills and piedmont in the Chihuahuan Desert. It extends into the Sky Island region to the west and the Edwards Plateau to the east. Substrates are typically derived from alluvium, often gravelly without a well-developed argillic or calcic soil horizon that would limit infiltration and storage of winter precipitation in deeper soil layers. *Prosopis* spp. and other deep-rooted shrubs exploit this deep soil moisture that is unavailable to grasses and cacti. Vegetation is typically dominated by *Prosopis glandulosa* or *Prosopis velutina* and succulents. Other desert scrub that may codominate or dominate includes *Acacia neovernicosa*, *Acacia constricta*, *Juniperus monosperma*, or *Juniperus coahuilensis*. Grass cover is typically low. During the last century, the area occupied by this system has increased through conversion of desert grasslands as a result of drought, overgrazing by livestock, and/or decreases in fire frequency. It is similar to Chihuahuan Mixed Desert and Thorn Scrub (CES302.734) but is generally found at higher elevations where *Larrea tridentata* and other desert scrub are not codominant. It is also similar to Chihuahuan Stabilized Coppice Dune and Sand Flat Scrub (CES302.737) but does not occur on eolian-deposited substrates.
- S059 *Colorado Plateau Blackbrush-Mormon Tea Shrubland* - This ecological system occurs in the Colorado Plateau on benchlands, colluvial slopes, pediments or bajadas. Elevation ranges from 560-1650 m. Substrates are shallow, typically calcareous, non-saline and gravelly or sandy soils over sandstone or limestone bedrock, caliche or limestone alluvium. It also occurs in deeper soils on sandy plains where it may have invaded desert grasslands. The vegetation is characterized by extensive open shrublands dominated by *Coleogyne ramosissima* often with *Ephedra viridis*, *Ephedra torreyana*, or *Grayia spinosa*. Sandy portions may include *Artemisia filifolia* as codominant. The herbaceous layer is sparse and composed of graminoids such as *Achnatherum hymenoides*, *Pleuraphis jamesii*, or *Sporobolus cryptandrus*.
- S061 *Chihuahuan Succulent Desert Scrub* - This ecological system is found in the Chihuahuan Desert on colluvial slopes, upper bajadas, sideslopes, ridges, canyons, hills and mesas. Sites are hot and dry. Gravel and rock are often abundant on the ground surface. The vegetation is characterized by the relatively high cover of succulent species such as *Agave lechuguilla*, *Euphorbia antisiphilitica*, *Fouquieria splendens*, *Ferocactus* spp., *Opuntia engelmannii*, *Opuntia imbricata*, *Opuntia spinosior*, *Yucca baccata*, and many others. Perennial grass cover is generally low. The abundance of succulents is diagnostic of this desert scrub system, but desert shrubs are usually present. This system does not include desert grasslands or shrub-steppe with a strong cacti component.
- S062 *Chihuahuan Mixed Desert and Thorn Scrub* - This widespread Chihuahuan Desert land cover type is composed of two ecological systems the Chihuahuan Creosotebush Xeric Basin Desert Scrub (CES302.731) and the Chihuahuan Mixed Desert and Thorn Scrub (CES302.734). This cover type includes xeric creosotebush basins and plains and the mixed desert scrub in the foothill transition zone above, sometimes extending up to the lower montane woodlands. Vegetation is characterized by *Larrea tridentata* alone or mixed with thornscrub and other desert scrub such as *Agave lechuguilla*, *Aloysia wrightii*, *Fouquieria splendens*, *Dasyliirion leiophyllum*, *Flourensia cernua*, *Leucophyllum minus*, *Mimosa aculeaticarpa* var. *biuncifera*, *Mortonia scabrella* (= *Mortonia sempervirens* ssp. *scabrella*), *Opuntia engelmannii*, *Parthenium incanum*, *Prosopis glandulosa*, and *Tiquilia greggii*. Stands of *Acacia constricta* *Acacia neovernicosa* or *Acacia greggii* dominated thornscrub are included in this system, and limestone substrates appear important for at least these species. Grasses such as *Dasyochloa pulchella*, *Bouteloua curtipendula*, *Bouteloua eriopoda*, *Bouteloua ramosa*, *Muhlenbergia porteri* and *Pleuraphis mutica* may be common, but generally have lower cover than shrubs.

Appendix D Cont.  
Code Description

- S063 *Sonoran Palo Verde-Mixed Cacti Desert Scrub* - This ecological system occurs on hillsides, mesas and upper bajadas in southern Arizona and extreme southeastern California. The vegetation is characterized by a diagnostic sparse, emergent tree layer of *Carnegia gigantea* (3-16 m tall) and/or a sparse to moderately dense canopy codominated by xeromorphic deciduous and evergreen tall shrubs *Parkinsonia microphylla* and *Larrea tridentata* with *Prosopis* sp., *Olneya tesota*, and *Fouquieria splendens* less prominent. Other common shrubs and dwarf-shrubs include *Acacia greggii*, *Ambrosia deltoidea*, *Ambrosia dumosa* (in drier sites), *Calliandra eriophylla*, *Jatropha cardiophylla*, *Krameria erecta*, *Lycium* spp., *Menodora scabra*, *Simmondsia chinensis*, and many cacti including *Ferocactus* spp., *Echinocereus* spp., and *Opuntia* spp. (both cholla and prickly pear). The sparse herbaceous layer is composed of perennial grasses and forbs with annuals seasonally present and occasionally abundant. On slopes, plants are often distributed in patches around rock outcrops where suitable habitat is present.
- S065 *Inter-Mountain Basins Mixed Salt Desert Scrub* - This extensive ecological system includes open-canopied shrublands of typically saline basins, alluvial slopes and plains across the Intermountain western U.S. This type also extends in limited distribution into the southern Great Plains. Substrates are often saline and calcareous, medium- to fine-textured, alkaline soils, but include some coarser-textured soils. The vegetation is characterized by a typically open to moderately dense shrubland composed of one or more *Atriplex* species such as *Atriplex confertifolia*, *Atriplex canescens*, *Atriplex polycarpa*, or *Atriplex spinifera*. Other shrubs present to codominate may include *Artemisia tridentata* ssp. *wyomingensis*, *Chrysothamnus viscidiflorus*, *Ericameria nauseosa*, *Ephedra nevadensis*, *Grayia spinosa*, *Krascheninnikovia lanata*, *Lycium* spp., *Picrothamnus desertorum*, or *Tetradymia* spp. *Sarcobatus vermiculatus* is generally absent, but if present does not codominate. The herbaceous layer varies from sparse to moderately dense and is dominated by perennial graminoids such as *Achnatherum hymenoides*, *Bouteloua gracilis*, *Elymus lanceolatus* ssp. *lanceolatus*, *Pascopyrum smithii*, *Pleuraphis jamesii*, *Pleuraphis rigida*, *Poa secunda*, or *Sporobolus airoides*. Various forbs are also present.
- S068 *Chihuahuan Stabilized Coppice Dune and Sand Flat Scrub* - This ecological system includes the open shrublands of vegetated coppice dunes and sandsheets found in the Chihuahuan Desert. Usually dominated by *Prosopis glandulosa* but includes *Atriplex canescens*, *Ephedra torreyana*, *Ephedra trifurca*, *Poliomintha incana*, and *Rhus microphylla* coppice sand scrub with 10-30% total vegetation cover. *Yucca elata*, *Gutierrezia sarothrae*, and *Sporobolus flexuosus* are commonly present.
- S069 *Sonora-Mojave Creosotebush-white Bursage Desert Scrub* - This ecological system forms the vegetation matrix in broad valleys, lower bajadas, plains and low hills in the Mojave and lower Sonoran deserts. This desert scrub is characterized by a sparse to moderately dense layer (2-50% cover) of xeromorphic microphyllous and broad-leaved shrubs. *Larrea tridentata* and *Ambrosia dumosa* are typically dominants, but many different shrubs, dwarf-shrubs, and cacti may codominate or form typically sparse understories. Associated species may include *Atriplex canescens*, *Atriplex hymenelytra*, *Encelia farinosa*, *Ephedra nevadensis*, *Fouquieria splendens*, *Lycium andersonii*, and *Opuntia basilaris*. The herbaceous layer is typically sparse, but may be seasonally abundant with ephemerals. Herbaceous species such as *Chamaesyce* spp., *Eriogonum inflatum*, *Dasyochloa pulchella*, *Aristida* spp., *Cryptantha* spp., *Nama* spp., and *Phacelia* spp. are common.
- S071 *Inter-Mountain Basins Montane Sagebrush Steppe* - This ecological system includes sagebrush communities occurring at montane and subalpine elevations across the western U.S. from 1000 m in eastern Oregon and Washington to over 3000 m in the southern Rockies. In British Columbia, it occurs between 450 and 1650 m in the southern Fraser Plateau and the Thompson and Okanagan basins. Climate is cool, semi-arid to subhumid. This system primarily occurs on deep-soiled to stony flats, ridges, nearly flat ridgetops, and mountain slopes. In general this system shows an affinity for mild topography, fine soils, and some source of subsurface moisture. It is composed primarily of *Artemisia tridentata* ssp. *vaseyana* (mountain sagebrush) and related taxa such as *Artemisia tridentata* ssp. *spiciformis* (= *Artemisia spiciformis*). *Purshia tridentata* may codominate or even dominate some stands. Other common shrubs include *Symphoricarpos* spp., *Amelanchier* spp., *Ericameria nauseosa*, *Peraphyllum ramosissimum*, *Ribes cereum*, and *Chrysothamnus*

Appendix D Cont.  
**Code Description**

---

- S071 Cont. viscidiflorus. Most stands have an abundant perennial herbaceous layer (over 25% cover), but this system also includes *Artemisia tridentata* ssp. *vaseyana* shrublands. Common graminoids include *Festuca arizonica*, *Festuca idahoensis*, *Hesperostipa comata*, *Poa fendleriana*, *Elymus trachycaulus*, *Bromus carinatus*, *Poa secunda*, *Leucopoa kingii*, *Deschampsia caespitosa*, *Calamagrostis rubescens*, and *Pseudoroegneria spicata*. In many areas, frequent wildfires maintain an open herbaceous-rich steppe condition, although at most sites, shrub cover can be unusually high for a steppe system (>40%), with the moisture providing equally high grass and forb cover.
- S074 *Southern Rocky Mountain Juniper Woodland and Savanna* - This ecological system occupies the lower and warmest elevations, growing from 1370 to 1830 m in a semi-arid climate, primarily along the east and south slopes of the southern Rockies and Arizona-New Mexico mountains. It is best represented just below the lower elevational range of ponderosa pine and often intermingles with grasslands and shrublands. This system is best described as a savanna that has widely spaced, mature (>150 years old) juniper trees and occasionally *Pinus edulis*. *Juniperus monosperma* and *Juniperus scopulorum* (at higher elevations) are the dominant tall shrubs or short trees. These savannas may have inclusions of more dense juniper woodlands and have expanded into adjacent grasslands during the last century. Graminoid species are similar to those found in Western Great Plains Shortgrass Prairie (CES303.672), with *Bouteloua gracilis* and *Pleuraphis jamesii* being most common. In addition, succulents such as species of *Yucca* and *Opuntia* are typically present.
- S075 *Inter-Mountain Basins Juniper Savanna* - This widespread ecological system occupies dry foothills and sandsheets of western Colorado, northwestern New Mexico, northern Arizona, Utah, west into the Great Basin of Nevada and southern Idaho. It is typically found at lower elevations ranging from 1500-2300 m. This system is generally found at lower elevations and more xeric sites than Great Basin Pinyon-Juniper Woodland (CES304.773) or Colorado Plateau Pinyon-Juniper Woodland (CES304.767). These occurrences are found on lower mountain slopes, hills, plateaus, basins and flats often where juniper is expanding into semi-desert grasslands and steppe. The vegetation is typically open savanna, although there may be inclusions of more dense juniper woodlands. This savanna is typically dominated by *Juniperus osteosperma* trees with high cover of perennial bunch grasses and forbs, with *Bouteloua gracilis*, *Hesperostipa comata*, and *Pleuraphis jamesii* being most common. In the southern Colorado Plateau, *Juniperus monosperma* or juniper hybrids may dominate the tree layer. Pinyon trees are typically not present because sites are outside the ecological or geographic range of *Pinus edulis* and *Pinus monophylla*.
- S077<sup>2</sup> *Chihuahuan Piedmont Semi-Desert Grassland* - This ecological system is a broadly defined desert grassland, mixed shrub-succulent or xeromorphic tree savanna that is typical of the Borderlands of Arizona, New Mexico and northern Mexico [Apacherian region] but extends west to the Sonoran Desert, north into the Mogollon Rim and throughout much of the Chihuahuan Desert. It is found on gently sloping bajadas that supported frequent fire throughout the Sky Islands and on mesas and steeper piedmont and foothill slopes in the Chihuahuan Desert. It is characterized by typically diverse perennial grasses. Common grass species include *Bouteloua eriopoda*, *Bouteloua hirsuta*, *Bouteloua rothrockii*, *Bouteloua curtipendula*, *Bouteloua gracilis*, *Eragrostis intermedia*, *Muhlenbergia porteri*, *Muhlenbergia setifolia*, *Pleuraphis jamesii*, *Pleuraphis mutica*, and *Sporobolus airoides*, succulent species of *Agave*, *Dasyllirion*, and *Yucca*, and tall-shrub/short-tree species of *Prosopis* and various oaks (e.g., *Quercus grisea*, *Quercus emoryi*, *Quercus arizonica*). Many of the historical desert grassland and savanna areas have been converted, some to Chihuahuan Mesquite Upland Scrub (CES302.733) (*Prosopis* spp.-dominated), through intensive grazing and other land uses.
- S079 *Inter-Mountain Basins Semi-Desert Shrub Steppe* - This widespread ecological system occurs throughout the intermountain western U.S. on dry plains and mesas, at approximately 1450 to 2320 m (4750-7610 feet) elevation. These grasslands occur in lowland and upland areas and may occupy swales, playas, mesatops, plateau parks, alluvial flats, and plains, but sites are typically xeric. Substrates are often well-drained sandy or loamy-textured soils derived from sedimentary parent materials but are quite variable and may include fine-textured soils derived from igneous and metamorphic rocks. When they occur near foothill grasslands they will be at lower elevations. The dominant perennial bunch grasses and shrubs within this system are all

Appendix D Cont.  
Code Description

- S079 very drought-resistant plants. These grasslands are typically dominated or codominated by *Achnatherum hymenoides*, *Aristida* spp., *Bouteloua gracilis*, *Hesperostipa comata*, *Muhlenbergia* sp., or *Pleuraphis jamesii* and may include scattered shrubs and dwarfshrubs of species of *Artemisia*, *Atriplex*, *Coleogyne*, *Ephedra*, *Gutierrezia*, or *Krascheninnikovia lanata*.
- S080 *Chihuahuan Gypsophilous Grassland and Steppe* - This ecological system is restricted to gypsum outcrops or sandy gypsiferous and/or often alkaline soils that occur in basins and slopes in the Chihuahuan Desert. Elevation range is from 1100-2000 m. These typically sparse grasslands, steppes or dwarf-shrublands are dominated by a variety of gypsophilous plants, many of which are endemic to these habitats. Characteristic species include *Tiquilia hispidissima*, *Atriplex canescens*, *Calylophus hartwegii*, *Ephedra torreyana*, *Frankenia jamesii*, *Bouteloua breviseta*, *Mentzelia perennis*, *Nama carnosum*, *Calylophus hartwegii* (= *Oenothera hartwegii*), *Selinocarpus lanceolatus*, *Sporobolus nealleyi*, *Sporobolus iroides*, and *Sartwellia flaveriae*. This system does not include the sparsely vegetated gypsum dunes that are included in North American Warm Desert Active and Stabilized Dune (CES302.744).
- S081 *Rocky Mountains Dry Tundra* - This widespread ecological system occurs above upper treeline throughout the Rocky Mountain cordillera, including alpine areas of ranges in Utah and Nevada, and isolated alpine sites in the northeastern Cascades. It is found on gentle to moderate slopes, flat ridges, valleys, and basins, where the soil has become relatively stabilized and the water supply is more or less constant. Vegetation in these areas is controlled by snow retention, wind desiccation, permafrost, and a short growing season. This system is characterized by a dense cover of low-growing, perennial graminoids and forbs. Rhizomatous, sod-forming sedges are the dominant graminoids, and prostrate and mat-forming plants with thick rootstocks or taproots characterize the forbs. Dominant species include *Artemisia arctica*, *Carex elynoides*, *Carex siccata*, *Carex scirpoidea*, *Carex nardina*, *Carex rupestris*, *Deschampsia caespitosa*, *Festuca brachyphylla*, *Festuca idahoensis*, *Geum rossii*, *Kobresia myosuroides*, *Phlox pulvinata*, and *Trifolium dasyphyllum*. Although alpine tundra dry meadow is the matrix of the alpine zone, it typically intermingles with alpine bedrock and scree, ice field, fell-field, alpine dwarfshrubland, and alpine/subalpine wet meadow systems.
- S083 *Rocky Mountain Subalpine Mesic Meadow* - This Rocky Mountain ecological system is restricted to sites in the subalpine zone where finely textured soils, snow deposition, or wind-swept dry conditions limit tree establishment. It is found typically above 3000 m in elevation in the southern part of its range and above 1500 m in the northern part. These upland communities occur on gentle to moderate gradient slopes. The soils are typically seasonally moist to saturated in the spring, but if so will dry out later in the growing season. These sites are not as wet as those found in Rocky Mountain Alpine-Montane Wet Meadow (CES306.812). Vegetation is typically forb-rich, with forbs contributing more to overall herbaceous cover than graminoids. Important taxa include *Erigeron* spp., *Asteraceae* spp., *Mertensia* spp., *Penstemon* spp., *Campanula* spp., *Lupinus* spp., *Solidago* spp., *Ligusticum* spp., *Thalictrum occidentale*, *Valeriana sitchensis*, *Balsamorhiza sagittata*, *Wyethia* spp., *Deschampsia caespitosa*, *Koeleria macrantha*, and *Dasiphora fruticosa*. Burrowing mammals can increase the forb diversity.
- S085 *Southern Rocky Mountain Montane Grassland* - This Rocky Mountain ecological system typically occurs between 2200 and 3000 m on flat to rolling plains and parks or on lower sideslopes that are dry, but it may extend up to 3350 m on warm aspects. Soils resemble prairie soils in that the Ahorizon is dark brown, relatively high in organic matter, slightly acid, and usually well-drained. An occurrence usually consists of a mosaic of two or three plant associations with one of the following dominant bunch grasses: *Danthonia intermedia*, *Danthonia parryi*, *Festuca idahoensis*, *Festuca arizonica*, *Festuca thurberi*, *Muhlenbergia filiculmis*, or *Pseudoroegneria spicata*. The subdominants include *Muhlenbergia montana*, *Bouteloua gracilis*, and *Poa secunda*. These large-patch grasslands are intermixed with matrix stands of spruce-fir, lodgepole, ponderosa pine, and aspen forests. In limited circumstances (e.g., South Park in Colorado), they form the "matrix" of high-elevation plateaus.
- S086 *Rocky Mountains Foothill Grassland* - This system typically occurs between 1600-2200 m in elevation. It is best characterized as a mixed-grass to tallgrass prairie on mostly moderate to gentle slopes, usually at the

Appendix D Cont.  
**Code Description**

---

- S086 Cont. base of foothill slopes, e.g., the hogbacks of the Rocky Mountain Front Range where it typically occurs as a relatively narrow elevational band between montane woodlands and shrublands and the shortgrass steppe, but extends east on the Front Range piedmont alongside the Chalk Bluffs along the Colorado-Wyoming border, out into the Great Plains on the Palmer Divide, and on piedmont slopes below mesas and foothills in northeastern New Mexico. A combination of increased precipitation from orographic rain, temperature, and soils limits this system to the lower elevation zone with approximately 40 cm of precipitation/year. It is maintained by frequent fire and associated with well-drained clay soils. Usually occurrences of this system have multiple plant associations that may be dominated by *Andropogon gerardii*, *Schizachyrium scoparium*, *Muhlenbergia montana*, *Nassella viridula*, *Pascopyrum smithii*, *Sporobolus cryptandrus*, *Bouteloua gracilis*, *Hesperostipa comata*, or *Hesperostipa neomexicana*. In Wyoming, typical grasses found in this system include *Pseudoroegneria spicata*, *Festuca idahoensis*, *Hesperostipa comata*, and species of *Poa*. Typical adjacent ecological systems include foothill shrublands, ponderosa pine savannas, juniper savannas, as well as shortgrass prairie.
- S088<sup>1</sup> *Western Great Plains Shortgrass Prairie* - This system is found primarily in the western half of the Western Great Plains Division in the rainshadow of the Rocky Mountains and ranges from the Nebraska Panhandle south into Texas and New Mexico, although grazing-impacted examples may reach as far north as southern Canada where it grades into Northwestern Great Plains Mixedgrass Prairie (CES303.674). This system occurs primarily on flat to rolling uplands with loamy, ustic soils ranging from sandy to clayey. In much of its range, this system forms the matrix system with *Bouteloua gracilis* dominating this system. Associated graminoids may include *Aristida purpurea*, *Bouteloua curtipendula*, *Bouteloua hirsuta*, *Buchloe dactyloides*, *Hesperostipa comata*, *Koeleria macrantha* (= *Koeleria cristata*), *Pascopyrum smithii* (= *Agropyron smithii*), *Pleuraphis jamesii*, *Sporobolus airoides*, and *Sporobolus cryptandrus*. Although mid-height grass species may be present, especially on more mesic land positions and soils, they are secondary in importance to the sod-forming short grasses. Sandy soils have higher cover of *Hesperostipa comata*, *Sporobolus cryptandrus*, and *Yucca elata*. Scattered shrub and dwarf-dwarf species such as *Artemisia filifolia*, *Artemisia frigida*, *Artemisia tridentata*, *Atriplex canescens*, *Eriogonum effusum*, *Gutierrezia sarothrae*, and *Lycium pallida* may also be present. Also, because this system spans a wide range, there can be some differences in the relative dominance of some species from north to south and from east to west. Large-scale processes such as climate, fire and grazing influence this system. High variation in amount and timing of annual precipitation impacts the relative cover of cool- and warm-season herbaceous species. In contrast to other prairie systems, fire is less important, especially in the western range of this system, because the often dry and xeric climate conditions can decrease the fuel load and thus the relative fire frequency within the system. However, historically, fires that did occur were often very expansive. Currently, fire suppression and more extensive grazing in the region have likely decreased the fire frequency even more, and it is unlikely that these processes could occur at a natural scale. A large part of the range for this system (especially in the east and near rivers) has been converted to agriculture. Areas of the central and western range have been impacted by the unsuccessful attempts to develop dryland cultivation during the Dust Bowl of the 1930s. The short grasses that dominate this system are extremely drought- and grazing-tolerant. These species evolved with drought and large herbivores and, because of their stature, are relatively resistant to overgrazing. This system in combination with the associated wetland systems represents one of the richest areas for mammals and birds. Endemic bird species to the shortgrass system may constitute one of the fastest declining bird populations.
- S090 *Inter-Mountain Basins Semi-Desert Grassland* - This widespread ecological system occurs throughout the intermountain western U.S. on dry plains and mesas, at approximately 1450 to 2320 m (4750-7610 feet) elevation. These grasslands occur in lowland and upland areas and may occupy swales, playas, mesatops, plateau parks, alluvial flats, and plains, but sites are typically xeric. Substrates are often well-drained sandy or loamy-textured soils derived from sedimentary parent materials but are quite variable and may include fine-textured soils derived from igneous and metamorphic rocks. When they occur near foothill grasslands they will be at lower elevations. The dominant perennial bunch grasses and shrubs within this system are all very drought-resistant plants. These grasslands are typically dominated or codominated by *Achnatherum hymenoides*, *Aristida* spp., *Bouteloua gracilis*, *Hesperostipa comata*, *Muhlenbergia* sp., or *Pleuraphis jamesii*

Appendix D Cont.  
Code Description

- S090 and may include scattered shrubs and dwarfshrubs of species of *Artemisia*, *Atriplex*, *Coleogyne*, *Ephedra*, *Gutierrezia*, or *Krascheninnikovia lanata*.
- S091<sup>4</sup> *Rocky Mountain Subalpine-Montane Riparian Shrubland* - This system is found throughout the Rocky Mountain cordillera from New Mexico north into Montana, and also occurs in mountainous areas of the Intermountain region and Colorado Plateau. These are montane to subalpine riparian shrublands occurring as narrow bands of shrubs lining streambanks and alluvial terraces in narrow to wide, low-gradient valley bottoms and floodplains with sinuous stream channels. Generally it is found at higher elevations, but can be found anywhere from 1700-3475 m. Occurrences can also be found around seeps, fens, and isolated springs on hillslopes away from valley bottoms. Many of the plant associations found within this system are associated with beaver activity. This system often occurs as a mosaic of multiple communities that are shrub- and herb-dominated and includes above-treeline, willow-dominated, snowmelt-fed basins that feed into streams. The dominant shrubs reflect the large elevational gradient and include *Alnus incana*, *Betula nana*, *Betula occidentalis*, *Cornus sericea*, *Salix bebbiana*, *Salix boothii*, *Salix brachycarpa*, *Salix drummondiana*, *Salix eriocephala*, *Salix geeyeriana*, *Salix monticola*, *Salix planifolia*, and *Salix wolfii*. Generally the upland vegetation surrounding these riparian systems are of either conifer or aspen forests.
- S092<sup>4</sup> *Rocky Mountain Subalpine-Montane Riparian Woodland* - This riparian woodland system is comprised of seasonally flooded forests and woodlands found at montane to subalpine elevations of the Rocky Mountain cordillera, from southern New Mexico north into Montana, and west into the Intermountain region and the Colorado Plateau. It occurs throughout the interior of British Columbia and the eastern slopes of the Cascade Mountains. This system contains the conifer and aspen woodlands that line montane streams. These are communities tolerant of periodic flooding and high water tables. Snowmelt moisture in this system may create shallow water tables or seeps for a portion of the growing season. Stands typically occur at elevations between 1500 and 3300 m (4920-10,830 feet), farther north elevation ranges between 900 and 2000 m. This is confined to specific riparian environments occurring on floodplains or terraces of rivers and streams, in V-shaped, narrow valleys and canyons (where there is cold-air drainage). Less frequently, occurrences are found in moderate-wide valley bottoms on large floodplains along broad, meandering rivers, and on pond or lake margins. Dominant tree species vary across the latitudinal range, although it usually includes *Abies lasiocarpa* and/or *Picea engelmannii*; other important species include *Pseudotsuga menziesii*, *Picea pungens*, *Picea engelmannii* X *glauca*, *Populus tremuloides*, and *Juniperus scopulorum*. Other trees possibly present but not usually dominant include *Alnus incana*, *Abies concolor*, *Abies grandis*, *Pinus contorta*, *Populus angustifolia*, *Populus balsamifera* ssp. *trichocarpa*, and *Juniperus osteosperma*.
- S093<sup>4</sup> *Rocky Mountain Lower Montane Riparian Woodland and Shrubland* - This system is found throughout the Rocky Mountain and Colorado Plateau regions within a broad elevation range from approximately 900 to 2800 m. This system often occurs as a mosaic of multiple communities that are tree-dominated with a diverse shrub component. This system is dependent on a natural hydrologic regime, especially annual to episodic flooding. Occurrences are found within the flood zone of rivers, on islands, sand or cobble bars, and immediate streambanks. They can form large, wide occurrences on mid-channel islands in larger rivers or narrow bands on small, rocky canyon tributaries and well-drained benches. It is also typically found in backwater channels and other perennially wet but less scoured sites, such as floodplains swales and irrigation ditches. Dominant trees may include *Acer negundo*, *Populus angustifolia*, *Populus balsamifera*, *Populus deltoides*, *Populus fremontii*, *Pseudotsuga menziesii*, *Picea pungens*, *Salix amygdaloides*, or *Juniperus scopulorum*. Dominant shrubs include *Acer glabrum*, *Alnus incana*, *Betula occidentalis*, *Cornus sericea*, *Crataegus rivularis*, *Forestiera pubescens*, *Prunus virginiana*, *Rhus trilobata*, *Salix monticola*, *Salix drummondiana*, *Salix exigua*, *Salix irrorata*, *Salix lucida*, *Shepherdia argentea*, or *Symphoricarpos* spp. Exotic trees of *Elaeagnus angustifolia* and *Tamarix* spp. are common in some stands. Generally, the upland vegetation surrounding this riparian system is different and ranges from grasslands to forests.
- S094<sup>4</sup> *North American Warm Desert Lower Montane Riparian Woodland and Shrubland* - This ecological system occurs in mountain canyons and valleys of southern Arizona, New Mexico, and adjacent Mexico and consists of mid- to low-elevation (1100-1800 m) riparian corridors along perennial and seasonally

Appendix D Cont.

**Code Description**

---

S094<sup>4</sup> intermittent streams. The vegetation is a mix of riparian woodlands and shrublands. Dominant trees include  
 Cont. *Populus angustifolia*, *Populus deltoides* ssp. *wislizeni*, *Populus fremontii*, *Platanus wrightii*, *Juglans major*,  
*Fraxinus velutina*, and *Sapindus saponaria*. Shrub dominants include *Salix exigua*, *Prunus* spp., *Alnus*  
*oblongifolia*, and *Baccharis salicifolia*. Vegetation is dependent upon annual or periodic flooding and  
 associated sediment scour and/or annual rise in the water table for growth and reproduction.

S095<sup>4</sup> *Western Great Plains Riparian Woodland and Shrubland* - This system is found in the riparian areas of  
 medium and small rivers and streams throughout the Western Great Plains. It is likely most common in the  
 Shortgrass Prairie and Northern Great Plains Steppe but extends west and as far as the Rio Grande in New  
 Mexico and into the Wyoming Basins in the north. It is found on alluvial soils in highly variable landscape  
 settings, from deep cut ravines to wide, braided streambeds. Hydrologically, these sites tended to be more  
 flashy with less developed floodplain than on larger rivers, and typically dried down completely for some  
 portion of the year. Dominant vegetation shares much with generally drier portions of larger floodplain  
 systems downstream, but overall abundance of vegetation is generally lower. Communities within this  
 system range from riparian forests and shrublands to gravel/sand flats. Dominant species include *Populus*  
*deltoides*, *Salix* spp., *Artemisia cana* ssp. *cana*, *Pascopyrum smithii*, *Sporobolus cryptandrus*, and  
*Schizachyrium scoparium*.

These areas are often subjected to heavy grazing and/or agriculture and can be heavily degraded. *Tamarix*  
 spp. and less desirable grasses and forbs can invade degraded examples up through central Colorado.  
 Another factor is that groundwater depletion and lack of fire have created additional species changes.  
 SWReGAP land cover mappers interpreted most of the riparian woodland and shrubland areas in the  
 Western Great Plains as this ecological system. Therefore, the SWReGAP map may include woody patches  
 of a similar landcover type, S120 Western Great Plains Floodplain, in this map class. The reverse may also  
 be true, where herbaceous patches of the S095 Western Great Plains Riparian Woodland and Shrubland  
 system may be mapped as S120.

S096<sup>4</sup> *Inter-Mountain Basins Greasewood Flat* - This ecological system occurs throughout much of the western  
 U.S. in Intermountain basins and extends onto the western Great Plains. It typically occurs near drainages on  
 stream terraces and flats or may form rings around more sparsely vegetated playas. Sites typically have  
 saline soils, a shallow water table and flood intermittently, but remain dry for most growing seasons. The  
 water table remains high enough to maintain vegetation, despite salt accumulations. This system usually  
 occurs as a mosaic of multiple communities, with open to moderately dense shrublands dominated or  
 codominated by *Sarcobatus vermiculatus*. *Atriplex canescens*, *Atriplex confertifolia*, or *Krascheninnikovia*  
*lanata* may be present to codominant. Occurrences are often surrounded by mixed salt desert scrub. The  
 herbaceous layer, if present, is usually dominated by graminoids. There may be inclusions of *Sporobolus*  
*airoides*, *Distichlis spicata* (where water remains ponded the longest), or *Eleocharis palustris* herbaceous  
 types.

S097<sup>4</sup> *North American Warm Desert Riparian Woodland and Shrubland* - This ecological system consists of low-  
 elevation (<1200 m) riparian corridors along medium to large perennial streams throughout canyons and the  
 desert valleys of the southwestern United States and adjacent Mexico. The vegetation is a mix of riparian  
 woodlands and shrublands. Dominant trees include *Acer negundo*, *Fraxinus velutina*, *Populus fremontii*,  
*Salix gooddingii*, *Salix lasiolepis*, *Celtis laevigata* var. *reticulata*, and *Juglans major*. Shrub dominants  
 include *Salix geyeriana*, *Shepherdia argentea*, and *Salix exigua*. Vegetation is dependent upon annual or  
 periodic flooding and associated sediment scour and/or annual rise in the water table for growth and  
 reproduction.

S098<sup>4</sup> *North American Warm Desert Riparian Mesquite Bosque* - This ecological system consists of low-elevation  
 (<1100 m) riparian corridors along intermittent streams in valleys of southern Arizona and New Mexico, and  
 adjacent Mexico. Dominant trees include *Prosopis glandulosa* and *Prosopis velutina*. Shrub dominants  
 include *Baccharis salicifolia*, *Pluchea sericea*, and *Salix exigua*. Vegetation, especially the mesquites, tap  
 groundwater below the streambed when surface flows stop. Vegetation is dependent upon annual rise in the  
 water table for growth and reproduction.

Appendix D Cont.  
Code Description

- S100<sup>4</sup> *North American Arid West Emergent Marsh* - This widespread ecological system occurs throughout much of the arid and semi-arid regions of western North America, typically surrounded by savanna, shrub steppe, steppe, or desert vegetation. Natural marshes may occur in depressions in the landscape (ponds, kettle ponds), as fringes around lakes, and along slow-flowing streams and rivers (such riparian marshes are also referred to as sloughs). Marshes are frequently or continually inundated, with water depths up to 2 m. Water levels may be stable, or may fluctuate 1 m or more over the course of the growing season. Water chemistry may include some alkaline or semi-alkaline situations, but the alkalinity is highly variable even within the same complex of wetlands. Marshes have distinctive soils that are typically mineral, but can also accumulate organic material. Soils have characteristics that result from long periods of anaerobic conditions in the soils (e.g., gleyed soils, high organic content, redoximorphic features). The vegetation is characterized by herbaceous plants that are adapted to saturated soil conditions. Common emergent and floating vegetation includes species of *Scirpus* and/or *Schoenoplectus*, *Typha*, *Juncus*, *Potamogeton*, *Polygonum*, *Nuphar*, and *Phalaris*. This system may also include areas of relatively deep water with floating-leaved plants (*Lemna*, *Potamogeton*, and *Brasenia*) and submergent and floating plants (*Myriophyllum*, *Ceratophyllum*, and *Elodea*).
- S102<sup>4</sup> *Rocky Mountain Alpine-Montane Wet Meadow* - These are high-elevation communities found throughout the Rocky Mountains and Intermountain regions, dominated by herbaceous species found on wetter sites with very low-velocity surface and subsurface flows. They range in elevation from montane to alpine (1000-3600 m). These types occur as large meadows in montane or subalpine valleys, as narrow strips bordering ponds, lakes, and streams, and along toeslope seeps. They are typically found on flat areas or gentle slopes, but may also occur on sub-irrigated sites with slopes up to 10%. In alpine regions, sites typically are small depressions located below late-melting snow patches or on snowbeds. Soils of this system may be mineral or organic. In either case, soils show typical hydric soil characteristics, including high organic content and/or low chroma and redoximorphic features. This system often occurs as a mosaic of several plant associations, often dominated by graminoids, including *Calamagrostis stricta*, *Caltha leptosepala*, *Cardamine cordifolia*, *Carex illota*, *Carex microptera*, *Carex nigricans*, *Carex scopulorum*, *Carex utriculata*, *Carex vernacula*, *Deschampsia caespitosa*, *Eleocharis quinqueflora*, *Juncus drummondii*, *Phippsia algida*, *Rorippa alpina*, *Senecio triangularis*, *Trifolium parryi*, and *Trollius laxus*. Often alpine dwarf-shrublands, especially those dominated by *Salix*, are immediately adjacent to the wet meadows. Wet meadows are tightly associated with snowmelt and typically not subjected to high disturbance events such as flooding.
- S108<sup>4</sup> *Western Great Plains Saline Depression* - This system is very similar to Northwestern Great Plains Open Freshwater Depression (CES303.675) and Western Great Plains Closed Depression Wetland (CES303.666). However, strongly saline soils cause both the shallow lakes and depressions and the surrounding areas to be more brackish. Salt encrustations can occur on the surface in some examples of this system, and the soils are severely affected and have poor structure. Species that typify this system are salt-tolerant and halophytic species such as *Distichlis spicata*, *Sporobolus airoides*, and *Hordeum jubatum*. During exceptionally wet years, an increase in precipitation can dilute the salt concentration in the soils of some of examples of this system which may allow for less salt-tolerant species to occur. Communities found within this system may also occur in floodplains (i.e., more open depressions), but probably should not be considered a separate system unless they transition to areas outside the immediate floodplain.
- S109<sup>2</sup> *Chihuahuan-Sonoran Desert Bottomland and Swale Grassland* - This ecological system occurs throughout the northern Chihuahuan Desert and adjacent Sky Islands and Sonoran Desert, as well as limited areas of the southern Great Plains and Edwards Plateau in relatively small depressions on broad mesas, plains and valley bottoms that receive runoff from adjacent areas. Water generally infiltrates relatively quickly. These depressions have deep, fine-textured soils that are neutral to slightly saline/alkaline. Vegetation is typically dominated by *Pleuraphis mutica* (tobosa swales) or other mesic graminoids such as *Pascopyrum smithii*, *Panicum obtusum*, *Sporobolus airoides*, or *Sporobolus wrightii*. With tobosa swales, sand-adapted species such as *Yucca elata* may grow at the swale's edge in the deep sandy alluvium that is deposited there from upland slopes. *Sporobolus airoides* and *Sporobolus wrightii* are more common in alkaline soils.

Appendix D Cont.  
**Code Description**

---

- S111<sup>3</sup> *Madrean Upper Montane Conifer-Oak Forest and Woodland* - This system occurs at the upper elevations in the Sierra Madre Occidentale and Sierra Madre Orientale. In the U.S., it is restricted to north and east aspects at high elevations (1980-2440 m) in the Sky Islands (Chiricahua, Huachuca, Pinaleno, Santa Catalina, and Santa Rita mountains) and along the Nantanes Rim. It is more common in Mexico and does not occur in Arizona central highlands. The vegetation is characterized by large- and small-patch forests and woodlands dominated by *Pseudotsuga menziesii*, *Abies coahuilensis*, or *Abies concolor* and Madrean oaks such as *Quercus hypoleucoides* and *Quercus rugosa*. It is similar to Rocky Mountain Montane Dry-Mesic Mixed Conifer Forest and Woodland (CES306.823).
- S112 *Madrean Pinyon-Juniper Woodland* - This system occurs on foothills, mountains and plateaus in the Sierra Madre Occidentale and Sierra Madre Orientale in Mexico, Trans-Pecos Texas, southern New Mexico and Arizona, generally south of the Mogollon Rim. Substrates are variable, but soils are generally dry and rocky. The presence of *Pinus cembroides*, *Pinus discolor*, or other Madrean trees and shrubs is diagnostic of this woodland system. *Juniperus coahuilensis*, *Juniperus deppeana*, *Juniperus pinchotii*, *Juniperus monosperma*, and/or *Pinus edulis* may be present to dominant. Madrean oaks such as *Quercus arizonica*, *Quercus emoryi*, *Quercus grisea*, or *Quercus mohriana* may be codominant. *Pinus ponderosa* is absent or sparse. If present, understory layers are variable and may be dominated by shrubs or graminoids.
- S113 *Chihuahuan Sandy Plains Semi-Desert Grassland* - This ecological system occurs across the Chihuahuan Desert and extends into the southern Great Plains where soils have a high sand content. These dry grasslands or steppe are found on sandy plains and sandstone mesas. The graminoid layer is dominated or codominated by *Achnatherum hymenoides*, *Bouteloua eriopoda*, *Bouteloua hirsuta*, *Hesperostipa neomexicana*, *Pleuraphis jamesii*, *Sporobolus cryptandrus*, *Sporobolus airoides*, or *Sporobolus flexuosus*. Typically, there are found scattered desert shrubs and stem succulents such as *Ephedra torreyana*, *Ephedra trifurca*, *Fallugia paradoxa*, *Prosopis glandulosa*, *Yucca elata*, and *Yucca torreyi* that are characteristic of the Chihuahuan Desert.
- S115 *Madrean Juniper Savanna* - This Madrean ecological system occurs in lower foothills and plains of southeastern Arizona, southern New Mexico extending into west Texas and Mexico. These savannas have widely spaced mature juniper trees and moderate to high cover of graminoids (>25% cover). The presence of Madrean *Juniperus* spp. such as *Juniperus coahuilensis*, *Juniperus pinchotii*, and/or *Juniperus deppeana* is diagnostic. *Juniperus monosperma* may be present in some stands, and *Juniperus deppeana* has a broader range than this Madrean system and extends north into southern stands of Southern Rocky Mountain Juniper Savanna and Woodland (CES306.834). Stands of *Juniperus pinchotii* may be short and resemble a shrubland. Graminoid species are a mix of those found in Western Great Plains Shortgrass Prairie (CES303.672) and Chihuahuan Piedmont Semi-Desert Grassland (CES302.735), with *Bouteloua gracilis* and *Pleuraphis jamesii* being most common. In addition, these areas include succulents such as species of *Yucca*, *Opuntia*, and Agave. Juniper savanna expansion into grasslands has been documented in the last century.
- S116 *Chihuahuan Mixed Salt Desert Scrub* - This system includes extensive open-canopied shrublands of typically saline basins in the Chihuahuan Desert. Stands often occur on alluvial flats and around playas. Substrates are generally fine-textured, saline soils. Vegetation is typically composed of one or more *Atriplex* species such as *Atriplex canescens*, *Atriplex obovata*, or *Atriplex polycarpa* along with species of *Allenrolfea*, *Flourensia*, *Salicornia*, *Suaeda*, or other halophytic plants. Graminoid species may include *Sporobolus airoides*, *Pleuraphis mutica*, or *Distichlis spicata* at varying densities.
- S117 *Coahuilan Chaparral* - This ecological system occurs in mountains across southeastern New Mexico (Guadalupe Mountains) and Trans-Pecos Texas (Chisos Mountains). It often dominates along the mid-elevation transition from the Chihuahuan Desert into mountains (1700-2500 m). It occurs on foothills, mountain slopes and canyons in drier habitats below the encinal and pine woodlands and is often associated with more xeric and coarse-textured substrates such as limestone, basalt or alluvium, especially in transition areas with more mesic woodlands. The moderate to dense shrub canopy includes many shrub oak species such as *Quercus intricata*, *Quercus pringlei*, *Quercus invaginata*, *Quercus laceyi*, *Quercus grisea*, *Quercus emoryi*, *Quercus toumeyii*, several widespread chaparral species such as *Arctostaphylos pungens*, *Ceanothus*

Appendix D Cont.  
Code Description

- S117 greggii, Fallugia paradoxa, and Garrya wrightii, and species characteristic of this system such as Arbutus arizonica, Arbutus xalapensis (= Arbutus texana), Fraxinus greggii, Fendlera rigida (= Fendlera linearis), Garrya ovata, Purshia mexicana (= ssp. mexicana), Rhus virens var. choriophylla (= Rhus choriophylla), and endemics Salvia lycioides (= Salvia ramosissima), Salvia roemeriana, and Salvia regla. Most chaparral species are fire-adapted, resprouting vigorously after burning or producing fire-resistant seeds. Stands occurring within montane woodlands are seral and a result of recent fires.
- S129 *Sonoran Mid-Elevation Desert Scrub* - This transitional desert scrub system occurs along the northern edge of the Sonoran Desert in an elevational band along the lower slopes of the Mogollon Rim/Central Highlands region between 750 and 1300 m. Stands occur in the Bradshaw, Hualapai, and Superstition mountains, among other desert ranges, and are found above Sonoran Paloverde-Mixed Cacti Desert Scrub (CES302.761) and below Mogollon Chaparral (CES302.741). Sites range from a narrow strip on steep slopes to very broad areas such as the Verde Valley. Climate is too dry for chaparral species to be abundant, and freezing temperatures during winter are too frequent and prolonged for many of the frost-sensitive species that are characteristic of Sonoran Paloverde-Mixed Cacti Desert Scrub (CES302.761), such as Carnegiea gigantea, Parkinsonia microphylla, Prosopis spp., Olneya tesota, Ferocactus sp., and Opuntia bigelovii. Substrates are generally rocky soils derived from parent materials such as limestone, granitic rocks or rhyolite. The vegetation is typically composed of an open shrub layer of Larrea tridentata, Ericameria linearifolia, or Eriogonum fasciculatum with taller shrub such as Canotia holacantha (limestone or granite) or Simmondsia chinensis (rhyolite). The herbaceous layer is generally sparse.
- S136 *Southern Colorado Plateau Sand Shrubland* - This large-patch ecological system is found on the south-central Colorado Plateau in northeastern Arizona extending into southern and central Utah. It occurs on windswept mesas, broad basins and plains at low to moderate elevations (1300-1800 m). Substrates are stabilized sandsheets or shallow to moderately deep sandy soils that may form small hummocks or small coppice dunes. This semi-arid, open shrubland is typically dominated by short shrubs (10-30 % cover) with a sparse graminoid layer. The woody layer is often a mixture of shrubs and dwarf-shrubs. Characteristic species include Ephedra cutleri, Ephedra torreyana, Ephedra viridis, and Artemisia filifolia. Coleogyne ramosissima is typically not present. Poliomintha incana, Parryella filifolia, Quercus havardii var. tuckeri, or Ericameria nauseosa may be present to dominant locally. Ephedra cutleri and Ephedra viridis often assume a distinctive matty growth form. Characteristic grasses include Achnatherum hymenoides, Bouteloua gracilis, Hesperostipa comata, and Pleuraphis jamesii. The general aspect of occurrences is an open low shrubland but may include small blowouts and dunes. Occasionally grasses may be moderately abundant locally and form a distinct layer. Disturbance may be important in maintaining the woody component. Eolian processes are evident, such as pediceled plants, occasional blowouts or small dunes, but the generally higher vegetative cover and less prominent geomorphic features distinguish this system from Inter-Mountain Basins Active and Stabilized Dune (CES304.775).
- S138 *Western Great Plains Mesquite Woodland and Shrubland* - This system is found primarily in the southern portion of the Western Great Plains Division, primarily in Texas, Oklahoma and eastern New Mexico. This system is dominated by Prosopis glandulosa with shortgrass species in the understory. Ziziphus obtusifolia and Atriplex canescens can codominate in some examples as can Opuntia species in heavily grazed areas. Historically this system probably occurred as a natural component on more fertile soils and along drainages.
- D02 *Recently Burned* - Areas that have burned in the recent past that are clearly evident in the imagery (images acquired between 1999-2001).
- D03 *Recently Mined or Quarried* - Areas where open pit mining or quarries are visible in the imagery (images acquired between 1999-2001), and are 2 hectares or greater in size.
- D04<sup>4</sup> *Invasive Southwest Riparian Woodland and Shrubland* - Areas that are dominated by introduced riparian woody species such as: Tamarix spp. and Elaeagnus angustifolius.

Appendix D Cont.

**Code Description**

---

- D06 *Invasive Perennial Grassland* - Areas that are dominated by introduced perennial grass species such as: *Agropyron cristatum*, *Bromus inermis*, *Eragrostis lehmannianna*, *Pennisetum spp.*, *Poa bulbosa*, *P. pratensis*, *Thinopyrum intermedium*.
- D09 *Invasive Annual and Biennial Forbland* - Areas that are dominated by introduced annual and/or biennial forb species such as: *Halogeton glomeratum*, *Kochia scoparia*, *Salsola spp.*, .
- D10 *Recently Logged Areas* - Areas that have recently been clear-cut or thinned by 50% or more and are clearly evident in the imagery (images acquired between 1999-2001).
- D11 *Recently Chained Pinyon-Juniper Areas* - Areas that have recently been chained to remove Pinyon-Juniper and are clearly evident in the imagery (images acquired between 1999-2001).
- N11 *Open water* - Areas of open water, generally with less than 25% cover of vegetation or soil.
- N21 *Developed, Low Intensity* - Open Space: Includes areas with a mixture of some construction materials, but mostly vegetation in the form of lawn grasses. Impervious surfaces account for less than 20 percent of total cover. These areas most commonly include largelot single-family housing units, parks, golf courses, and vegetation planted in developed settings for recreation, erosion control, or aesthetic purposes. Developed, Low Intensity: Includes areas with a mixture of constructed materials and vegetation. Impervious surfaces account for 20-49 percent of total cover. These areas most commonly include singlefamily housing units.
- N22 *Developed, Medium - High Intensity* - Developed, Medium Intensity: Includes areas with a mixture of constructed materials and vegetation. Impervious surface accounts for 50-79 percent of the total cover. These areas most commonly include single-family housing units. Developed, High Intensity: Includes highly developed areas where people reside or work in high numbers. Examples include apartment complexes, row houses and commercial/industrial. Impervious surfaces account for 80 to 100 percent of the total cover.
- N31 *Barren, Non-Specific* - (Rock/Sand/Clay)-Barren areas of bedrock, desert pavement, scarps, talus, slides, volcanic material, glacial debris, sand dunes, strip mines, gravel pits and other accumulation of earthen material. Generally, vegetation accounts for less than 15% of total cover.
- N80 *Agriculture* - An aggregated landcover type that includes both Pasture/Hay (N81): areas of grasses, legumes, or grass-legume mixtures planted for livestock grazing or the production of seed or hay crops, typically on a perennial cycle, where pasture/hay vegetation accounts for greater than 20 percent of total vegetation, and Cultivated Crops (N82): areas used for the production of annual crops, such as corn, soybeans, vegetables, tobacco, and cotton, and also perennial woody crops such as orchards and vineyards, where crop vegetation accounts for greater than 20 percent of total vegetation. N82 also includes all land being actively tilled.

---

<sup>1</sup> Priority habitats for New Mexico CWCS.

<sup>2</sup> Aggregated SWReGAP land cover types into Chihuahuan Semi-Desert Grassland priority habitat.

<sup>3</sup> Aggregated SWReGAP land cover types into Madrean Pine-Oak/Conifer-Oak Forest and Woodland priority habitat.

<sup>4</sup> Aggregated SWReGAP land cover types into Riparian priority habitat.

<sup>5</sup> Aggregated SWReGAP land cover types into Rocky Mountain Montane Mixed Conifer Forest and Woodland priority habitat.